

READ LABEL BEFORE USE

WARNING: Keep out of reach of children

KIWICARE[®]

Weed Weapon Extra Strength Commercial Concentrate 20 L

Post-emergent herbicide with rapid activity for effective control of annual and perennial grasses and weeds

Developed and tested for commercial use in New Zealand conditions

Proudly New Zealand made

CONTAINS: 285.1 g/L Glyphosate and 5.6 g/L Saflufenacil in the form of a soluble concentrate

HSNO Approval Code: HSR100724

HSNO Classifications: 6.3B, 6.4A, 6.8B, 9.1B

Kiwicare Corporation Limited, 225 Maces Road,
Christchurch, New Zealand
Ph: 64 3 389 0778



READ LABEL BEFORE USE

- Kills a broad range of weeds and grasses to the roots
- Can be used all year round
- Ideal for hard to kill weeds in garden areas, parks, reserves, roadsides, paths driveways, flower beds and other non-crop area (avoid contact with desirable plants)
- Biodegradable in soil
- Rainfast in 2 hours
- Replant after 5 days

DIRECTIONS

Post emergent weed control

THOROUGHLY SHAKE OR AGITATE CONCENTRATE PRIOR TO DECANTING. Apply to weed foliage to the point of run off.

Beware: Apply this product carefully; spray drift may cause serious damage to desirable plants.

Application Methods

Knapsack and Handgun Application: Use conventional application equipment capable of achieving complete coverage of target plants.

Application Rates

Environment <i>Not registered for use in or around food crops</i>	Dilution Rates		Critical Comments
	Knapsack ml/10 L	Handgun L/100 L	
Annual Weeds	125	1.25	Apply at higher rate for larger and hard to kill weeds. Ensure full coverage of weed foliage. For optimum results, apply in warm, sunny weather when weeds are actively growing.
Perennial Weeds	125 - 250	1.25 - 2.5	
Woody Weeds	125 - 250	1.50 - 2.50	
Creeping Weeds e.g. Tradescantia, ivy, periwinkle	125	1.25	Ensure entire 'colony' is treated including lower levels of foliage. For best results on waxy leaved plants e.g. ivy, add a suitable wetting agent.

For Best Results

- Apply in dry conditions when rain is not expected for 2 hours
- Complete coverage is required for effective control
- Avoid contact with foliage, new bark or near roots of desirable plants. If sprayed, immediately hose off with water
- Best applied in temperature above 10°C
- Can be used with a spray marker dye
- Wait 5 days before planting in sprayed areas

WARNING: May cause skin and eye irritation. Suspected of damaging fertility or the unborn child. Toxic to aquatic life with long-lasting effects.

Precautions: Wear waterproof gloves and eye protection when mixing and spraying, and wash hands and exposed skin thoroughly after use. Avoid skin contact and inhalation of spray. Rinse equipment thoroughly after use and dispose of surplus spray and washings well away from susceptible plants. Do not use around fruit vines, avocados or sensitive plants.

First Aid: If swallowed rinse mouth and drink 1 - 2 glasses of water or milk. Do not induce vomiting. If eye contact occurs, flush eyes with running water for several minutes. Remove contact lenses if present. If spilled on skin, wash with soap and water. If skin or eye irritation persists, seek medical advice or contact the National Poisons Centre on 0800 764 766 and have product label at hand.

Environmental Care: This product is toxic to aquatic environments. Collect any spillage. Do not use on or around food or animal feed producing plants. This substance must not be applied in a wide-dispersive manner into, onto or over water.

Storage/Disposal: Store in original container in a cool dark place away from children and foodstuffs. Triple rinse and dispose of container in the rubbish.

As Kiwicare Corporation Limited has no control over the storage or use of this product, no liability is accepted for accidents or damage arising from the incorrect storage or use of this product.

